

# Patient case studies: Navigating challenging clinical scenarios

Professor Lorient

Gustave Roussy Cancer Institute Université Paris-Saclay, Paris, France

**Prescribing Information is available at the end of this presentation.**

This promotional meeting is fully sponsored and supported by Astellas, including speaker-related honoraria and production of materials. It is intended for healthcare professionals only.

EV, in combination with P, is indicated for the 1L treatment of adult patients with unresectable/mUC who are eligible for platinum-containing chemotherapy.<sup>1</sup>

Please note: This indication has received EMA approval; reimbursement in some EU countries is still pending.

EV as monotherapy is indicated for the treatment of adult patients with LA/mUC who have previously received a platinum-containing chemotherapy and a PD-1/L1 inhibitor.<sup>1</sup>

1L, first line; AE, adverse event; EMA, European Medicines Agency; EU, European Union;

EV, enfortumab vedotin; LA, locally advanced; mUC, metastatic urothelial carcinoma;

P, pembrolizumab; PD-1/L1, programmed death-1/ligand-1.

1. PADCEV™ (enfortumab vedotin). Summary of Product Characteristics.

Date of preparation: July 2025 | Job code: MAT-NL-PAD-2025-00037



This medicinal product is subject to additional monitoring.

**NL: Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system. Nederland:

Nederlands Bijwerkingen Centrum Lareb;

Website: [www.lareb.nl](http://www.lareb.nl)

**UK: Adverse events should be reported.**

Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for 'MHRA yellow card' in the Google Play Store or Apple App Store.

Adverse events should also be reported to Astellas Pharma Ltd on 0800 783 5018



**PADCEV™**  
enfortumab vedotin  
Injection for IV infusion 20 mg & 30 mg vials



astellas

# Speaker disclosures

- **Consultancy:** AstraZeneca, Bristol Myers Squibb, Clovis Oncology, Janssen, Merck KGaA, Merck Sharp & Dohme, Nektar, Taiho, Pfizer, Roche, Seattle Genetics
- **Research funding:** Celsius, Janssen, Merck Sharp & Dohme, Roche, Sanofi
- **Speaking activities:** Astellas Pharma, Bristol Myers Squibb, Janssen, Merck Sharp & Dohme, Pfizer
- **Travel support:** AstraZeneca, Bristol Myers Squibb, Janssen, Merck Sharp & Dohme, Seattle Genetics
- **Clinical study investigator:** Astellas Pharma, AstraZeneca, Basilea, Gilead, Bristol Myers Squibb, Clovis Oncology, Incyte, Janssen, Merck KGaA, Merck Sharp & Dohme, Nektar, Pfizer, Roche, Seattle Genetics

# Rare histological subtypes/mixed histology in advanced UC



Clinical  
scenario 3

# Poll: Vote via the Onomi app!




**To what extent do variant histologies influence your treatment decisions in advanced UC?**

- Significantly
- Moderately
- Not at all
- Unsure/depends on the variant



**Vote via the  
Onomi app**

# Clinical guidelines are suited for tumours with UC-predominant histology



The ESMO Clinical Practice Guideline aligns with the EU regulatory approval for pembrolizumab 'for the 1L treatment of unresectable or metastatic UC in adults'<sup>1,2</sup>


Figure adapted from Powles T et al. 2024.

\*FDA approved and EMA approved;<sup>1,2</sup> †Rechallenge with single-agent ICI is not encouraged without further evidence;<sup>1</sup> ‡In tumours with selected FGFR DNA fusions and mutations;<sup>1</sup> ¶ESMO-MCBS v1.1 was used to calculate scores for new therapies/indications approved by the EMA or FDA. The scores have been calculated and validated by the ESMO-MCBS Working Group and reviewed by the authors (<https://www.esmo.org/guidelines/esmo-mcbs/esmo-mcbs-evaluation-forms>);<sup>1</sup> §This should be assessed within 10 weeks of completion of chemotherapy;<sup>1</sup> ††Rechallenge with PBCT may be considered if progression occurred 12 months after the end of previous PBCT or 12 months after the end of previous PBCT and maintenance avelumab;<sup>1</sup> †††Platinum doublets to be considered if the treatment-free interval from the last PBCT is >1 year;<sup>1</sup> ¶¶To be considered when other therapies are not available.<sup>1</sup>

1L, first line; Carbo; carboplatin; Cis, cisplatin; EMA, European Medicines Agency; ESMO, European Society for Medical Oncology; EV, enfortumab vedotin; FDA, US Food and Drug Administration; FGFR, fibroblast growth factor receptor; Gem, gemcitabine; ICI, immune checkpoint inhibitor; m, metastatic; MCBS, Magnitude of Clinical Benefit Scale; P, pembrolizumab; PBCT, platinum-based chemotherapy; UC, urothelial carcinoma.

1. Powles T et al. *Ann Oncol* 2024;35:485–490; 2. EMA. Summary of opinion. July 2024. Available at: [ema.europa.eu/en/documents/smop/chmp-post-authorisation-summary-positive-opinion-padcev-ii-13\\_en.pdf](https://ema.europa.eu/en/documents/smop/chmp-post-authorisation-summary-positive-opinion-padcev-ii-13_en.pdf). Last accessed: June 2025.


# Clinical guidelines are suited for tumours with predominant UC histology



The ESMO Clinical Practice Guideline aligns with the EU regulatory approval for pembrolizumab ‘for the 1L treatment of unresectable or metastatic UC in adults’

# Most common variant histologies in bladder cancer

Distribution of histologic subtypes of bladder cancer in the study population within the National Cancer Database (n=4469)




■ Squamous      ■ Adenocarcinoma      ■ Neuroendocrine  
■ Sarcomatoid      ■ Micropapillary      ■ Plasmacytoid

# Neuroendocrine subtype

## Key features:

- Aggressive tumour, but it often responds well to chemotherapy<sup>1</sup>
- Treatment for this histologic variant has been extrapolated from data on small cell lung cancer<sup>2</sup>
- Clinical trials are currently investigating the combinations of chemotherapy + IO or lurbinectedin + avelumab<sup>3,4</sup>




**Neuroendocrine subtype<sup>1</sup>**



# Sarcomatoid subtype

## Key features:

- Characterised by malignant spindle cells with a non-specific morphologic appearance<sup>1</sup>
- Conflicting data about the role of neoadjuvant chemotherapy<sup>1</sup>
- High PD-L1 expression and data suggesting activity of IO<sup>2</sup>




**Sarcomatoid subtype<sup>1</sup>**

# Squamous cell carcinoma subtype

## Key features:

- Keratinisation and intracellular bridges<sup>1,2</sup>
- Aggressive; frequently presents at locally advanced stage<sup>2</sup>
- Lower response rate to chemotherapy (vs. cystectomy)<sup>1</sup>




**Squamous cell carcinoma subtype<sup>1</sup>**

# Micropapillary subtype

## Key features:

- Small nests of tumour cells within lacunae and without fibrovascular cores
- More likely to be diagnosed at an advanced stage
- HER2 overexpression



**Micropapillary subtype**

# Patient profile: Jean



Jean

- Male
- **Age:** 69 years
- **ECOG PS:** 1
- **GFR:** 67 ml/min
- **BMI:** 23 kg/m<sup>2</sup>
- **HbA1c:** 6.3%



**Lifestyle:** Former tobacco exposure  
**Employment status/job:** Retired, former fireman  
**Family history of cancer:** None



**Comorbidities:** Moderate hypertension



**Concomitant medications:** Enalapril



**Any other relevant medical history:** NA

**Habits:** Physically active and volunteers in a non-profit organisation



# Patient treatment/disease history



- Prior surgery/treatments: Diagnosis of MIBC with 30% UC and 70% of SCC
- CT scan showed initial enlarged pelvic LNs and in LN in the retroperitoneum
- Confirmation of mixed histology after second pathological assessments
- PS 1, creatinine clearance of 70 ml/min
- At this time: SOC was PBCT +/- avelumab
- Three cycles of CisGem was given and then three cycles of CarboGem (moderate renal failure); SD was the best response
- After discussion with the patient, avelumab was initiated
- New disease progression
- Liquid biopsy analysis found: *FGFR3* R248C (2.2%), S249C (0.1%), *MDM2* amplification, *CDKN2A* deletion


# Potential treatment options

Erdafitinib or clinical trial testing specific FGFR3 inhibitor

EV


Clinical trial targeting PRMT5

Clinical trial with other ADC (e.g., TROP2)



**After discussing with the patient, we started EV at standard dosing (1.25 mg/kg)**

## After 3 months: Moderate PD



Patient was switched to a pan-FGFR inhibitor with clinical progression

# Current data on EV in variant subtypes

Publication	Population	Total ORR (%)	VH response
Phase II	Pure UC 84, UC/Sq 15, UC/VH 26	44	Not reported
Phase I	N=155	43	Not reported
Phase III	N=301 EV 229 UC / 45 with VH	40.6	Not reported
Post-CPI	62 UC, 12 UC/Sq, 15 UC/VH	52	Not reported
Phase II EV+P Cis-ineligible EV-103 Cohort A	15 UC, 8 UC/Sq, 22 UC/VH	73	Not reported
EV +/-P Cis-ineligible EV-103 Cohort K	76 EV+P 73 EV	64.5 (EV+P) 45.2 (EV)	Not reported
Phase III 1L EV-302	379 UC, 50 UC/VH, four pure VH	67.7 (EV+P)	Not reported

***Table is for illustrative purposes only; Direct comparisons should not be made***

Cis, cisplatin; CPI, checkpoint inhibitor; EV, enfortumab vedotin; ORR, overall response rate; P, pembrolizumab; Sq, squamous; UC, urothelial carcinoma; VH, variant histology.  
 UroToday. ASCO 2024: What Is Variant Histology Urothelial Cancer and What Are the Available Treatment Options? Available at: <https://www.urotoday.com/conference-highlights/asco-2024/asco-2024-bladder-cancer/152654-asco-2024-what-is-variant-histology-urothelial-cancer-and-what-are-the-available-treatment-options.html>. Last accessed: July 2025.



# Current data on EV in variant subtypes

**Analysis of the UNITE RWE study:** Multi-site retrospective study of patients with advanced UC treated with targeted agents, including EV

Variant	ORR (%)	UC predominant (<50% HV), n	ORR (%)	HV predominant (50–99% HV), n	ORR (%)	pHV (100% HV), n	ORR (%)
Squamous (n=94)	47 (36/76)	70	55 (31/56)	17	33 (5/15)	7	0 (0/5)
Micropapillary (n=41)	35 (12/34)	35	38 (11/29)	6	20 (1/5)	0	-
Plasmacytoid (n=23)	53 (9/17)	18	64 (9/14)	2	NE	3	0 (0/3)
Sarcomatoid (n=21)	47 (8/17)	15	38 (5/13)	4	100 (3/3)	2	0 (0/1)
Adenocarcinoma/glandular (n=9)	56 (5/9)	8	63 (5/8)	1	0 (0/1)	0	-
NE/small cell (n=9)	0 (0/8)	3	0 (0/3)	4	0 (0/3)	2	0 (0/2)
Nested (n=2)	50 (1/2)	1	0 (0/1)	1	100 (1/1)	0	-
Lipid cell variant (n=1)	100 (1/1)	1	100 (1/1)	0	-	0	-
Any VH	44 (72/164)	151	50 (62/125)	35	36 (10/28)	14	0 (0/11)

# Current data on EV+P in variant subtypes

**Single-centre retrospective cohort study:** Patients with mUC who received 1L treatment with EV+P\*

	Transitional UC predominant (n=37)	Transitional UC <50% VH (squamous/glandular) (n=6)	Transitional UC >50% VH (squamous/glandular) (n=6)	Micropapillary variant (n=7)	Other VHs (n=8)
ORR (%)	57.6 (15/26)	16.6 (1/6)	0 (0/11)	57.1 (4/7)	25 (2/8)
PFS (12 m) (%)	45.9 (17/37)	NA	30.7 (4/13)	42.8 (3/7)	12.5 (1/8)
OS (12 m) (%)	56.7 (21/37)	NA	38.4 (5/13)	57.1 (4/7)	37.5 (3/8)


Patients with transitional UC or other non-squamous/glandular histologies were significantly more likely to respond to treatment compared with those with any degree of squamous or glandular differentiation (p=0.001).

\*Between 1 June 2023, and 8 August 2024.


1L, first-line; EV, enfortumab vedotin; mUC, metastatic urothelial carcinoma; NA, not available; ORR, objective response rate; OS, overall survival; P, pembrolizumab; PFS, progression-free survival; UC, urothelial carcinoma; VH, variant histology. UroToday. ASCO 2025: Impact of Histological Subtypes in Patients Receiving First-Line Treatment with Enfortumab Vedotin/pembrolizumab in Advanced Metastatic Urothelial Cancer. Available at: <https://www.urotoday.com/conference-highlights/asco-2025/asco-2025-bladder-cancer/160989-asco-2025-impact-of-histological-subtypes-in-patients-receiving-first-line-treatment-with-enfortumab-vedotin-pembrolizumab-in-advanced-metastatic-urothelial-cancer.html?tmpl=component&print=1>. Last accessed: July 2025.

# Nectin-4 expression in molecular subtypes


Expression of Nectin-4 according to subtype in MIBC/mUC (N=98)<sup>1</sup>



Expression of Nectin-4 across the consensus molecular subtypes of MIBC and association with luminal markers<sup>2</sup>



Nectin-4 expression (H-score) in different variant histologies of bladder cancer<sup>3</sup>




MIBC, muscle-invasive bladder cancer; mUC, metastatic urothelial carcinoma; ns, not significant.

1. Lorient Y et al. *Nat Commun* 2024;15:8603 (suppl); 2. Chu CE et al. *Clin Cancer Res* 2022;27:5123–5130; 3. Rodler S et al. *Cancers* 2022;14:4411.

# Expression differences of Nectin-4, TROP-2, and HER2 in UC

Expression differences of Nectin-4, TROP-2, and HER2 4 in conventional UC and histologic variants



# Summary



Non-urothelial bladder cancer subtypes are uncommon and present a substantial challenge for clinicians, primarily because of the **scarcity of prospective data and their frequent exclusion from conventional UC trials**<sup>1</sup>



Each non-urothelial subtype exhibits **distinct histological and clinical characteristics** that often lead to more **aggressive courses and poorer prognoses than conventional UC**<sup>2</sup>



There is an **emerging role of immunotherapy and ADCs** with variable efficacy in different histological subtypes<sup>2–4</sup>



There is a critical need for **inclusive clinical trials and subtype-specific analyses**.<sup>2</sup> Analyses of the impact of variant histologies should be conducted in Phase III trials at a minimum<sup>5</sup>



Patients should be referred to specialised centres that offer **pathological review, molecular profiling and membrane protein profiling**<sup>5</sup>

ADC, antibody–drug conjugate; UC, urothelial carcinoma.

1. Vlachou E et al. *Am Soc Clin Oncol Educ Book* 2024;44:e438640; 2. Alderson M et al. *Bladder Cancer* 2020;6:107–122; 3. UroToday. ASCO 2024: What Is Variant Histology Urothelial Cancer and What Are the Available Treatment Options? Available at: <https://www.urotoday.com/conference-highlights/asco-2024/asco-2024-bladder-cancer/152654-asco-2024-what-is-variant-histology-urothelial-cancer-and-what-are-the-available-treatment-options.html>. Last accessed: July 2025; 4. Kwon W-A et al. *Biomedicines* 2025;13:86; 5. Speaker's own opinion.



Please refer to the EMA SmPC for  
PADCEV™ (enfortumab vedotin)  
via the following link:

[https://www.ema.europa.eu/en/documents/product-information/padcev-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/padcev-epar-product-information_en.pdf)

PADCEV is subject to medicinal prescription.  
Astellas Pharma B.V., Sylviusweg 62, 2333 BE Leiden, The Netherlands



Please scan the QR  
code to access the  
UK aPI for PADCEV



Please scan the QR  
code to access the NL  
SmPC for PADCEV



## ABBREVIATED SUMMARY OF PRODUCT CHARACTERISTICS

For full prescribing information refer to the Summary of Product Characteristics (SPC).

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. **NAME OF THE MEDICINAL PRODUCT:** Padcev 20 mg powder for concentrate for solution for infusion & Padcev 30 mg powder for concentrate for solution for infusion **QUALITATIVE AND QUANTITATIVE COMPOSITION:** Padcev 20 mg powder for concentrate for solution for infusion: One vial of powder for concentrate for solution for infusion contains 20 mg enfortumab vedotin. Padcev 30 mg powder for concentrate for solution for infusion: One vial of powder for concentrate for solution for infusion contains 30 mg enfortumab vedotin. After reconstitution, each mL of solution contains 10 mg of enfortumab vedotin. Enfortumab vedotin is comprised of a fully human IgG1 kappa antibody, conjugated to the microtubule-disrupting agent monomethyl auristatin E (MMAE) via a protease-cleavable maleimidecaproyl valine-citrulline linker. For the full list of excipients, see section 6.1 of the SPC.

**PHARMACEUTICAL FORM:** Powder for concentrate for solution for infusion. While to off-white lyophilized powder. **CLINICAL PARTICULARS: Therapeutic indications:** Padcev, in combination with pembrolizumab, is indicated for the first-line treatment of adult patients with unresectable or metastatic urothelial cancer who are eligible for platinum-containing chemotherapy. Padcev as monotherapy is indicated for the treatment of adult patients with locally advanced or metastatic urothelial cancer who have previously received a platinum-containing chemotherapy and a programmed death receptor-1 or programmed death-ligand 1 inhibitor (see section 5.1 of the SPC). **Posology and method of administration:** Treatment with Padcev should be initiated and supervised by a physician experienced in the use of anti-cancer therapies. Ensure good venous access prior to starting treatment (see section 4.4 of the SPC). **Posology:** As monotherapy, the recommended dose of enfortumab vedotin is 1.25 mg/kg (up to a maximum of 125 mg for patients >100 kg) administered as an intravenous infusion over 30 minutes on Days 1, 8 and 15 of a 28-day cycle until disease progression or unacceptable toxicity. When given in combination with pembrolizumab, the recommended dose of enfortumab vedotin is 1.25 mg/kg (up to a maximum of 125 mg for patients >100 kg) administered as an intravenous infusion over 30 minutes on Days 1 and 8 of every 3-week (21-day) cycle until disease progression or unacceptable toxicity. The recommended dose of pembrolizumab is either 200 mg every 3 weeks or 400 mg every 6 weeks administered as an intravenous infusion over 30 minutes. Patients should be administered pembrolizumab after enfortumab vedotin when given on the same day. Refer to the pembrolizumab SPC for additional dosing information of pembrolizumab.

**Table 1. Recommended dose reductions of enfortumab vedotin for adverse reactions**

	Dose level
Starting dose	1.25 mg/kg up to 125 mg
First dose reduction	1.0 mg/kg up to 100 mg
Second dose reduction	0.75 mg/kg up to 75 mg
Third dose reduction	0.5 mg/kg up to 50 mg

### Dose modifications

**Table 2. Dose interruption, reduction and discontinuation of enfortumab vedotin in patients with locally advanced or metastatic urothelial cancer**

Adverse reaction	Severity*	Dose modification*
	Suspected Stevens-Johnson syndrome (SJS) or toxic epidermal necrolysis (TEN) or bullous lesions	Immediately withhold and refer to specialised care.
<b>Skin reactions</b>	Confirmed SJS or TEN; Grade 4 or recurrent Grade 3	Permanently discontinue.
	Grade 2 worsening Grade 2 with fever Grade 3	<ul style="list-style-type: none"><li>Withhold until Grade ≤1.</li><li>Referral to specialised care should be considered.</li><li>Resume at the same dose level or consider dose reduction by one dose level (see Table 1).</li></ul>
<b>Hyperglycaemia</b>	Blood glucose >13.9 mmol/L (>250 mg/dL)	<ul style="list-style-type: none"><li>Withhold until elevated blood glucose has improved to ≤13.9 mmol/L (&lt;250 mg/dL).</li><li>Resume treatment at the same dose level.</li></ul>
<b>Pneumonitis/interstitial lung disease (ILD)</b>	Grade 2	<ul style="list-style-type: none"><li>Withhold until Grade ≤1, then resume at the same dose or consider dose reduction by one dose level (see Table 1).</li></ul>
	Grade ≥3	Permanently discontinue.
<b>Peripheral neuropathy</b>	Grade 2	<ul style="list-style-type: none"><li>Withhold until Grade ≤1.</li><li>For first occurrence, resume treatment at the same dose level.</li><li>For a recurrence, withhold until Grade ≤1, then resume treatment reduced by one dose level (see Table 1).</li></ul>
	Grade ≥3	Permanently discontinue.

\*Toxicity was graded per National Cancer Institute Common Terminology Criteria for Adverse Events Version 5.0 (NCI-CTCAE v5.0) where Grade 1 is mild, Grade 2 is moderate, Grade 3 is severe and Grade 4 is life threatening.

**Special populations: Elderly:** No dose adjustment is necessary in patients >65 years of age. **Renal impairment:** No dose adjustment is necessary in patients with mild [creatinine clearance (CrCL) ~60–90 mL/min], moderate (CrCL 30–60 mL/min) or severe (CrCL 15–30 mL/min) renal impairment. Enfortumab vedotin has not been evaluated in patients with end stage renal disease (CrCL <15 mL/min) (see section 5.2 of the SPC). **Hepatic impairment:** No dose adjustment is necessary in patients with mild hepatic impairment [total bilirubin of 1 to 1.5 × upper limit of normal (ULN) and AST and/or total bilirubin ≤ ULN and AST > ULN]. Enfortumab vedotin has only been evaluated in a limited number of patients with moderate and severe hepatic impairment. Hepatic impairment is expected to increase the systemic exposure to MMAE (the cytotoxic drug); therefore, patients should be closely monitored for potential adverse events. Due to the sparsity of the data in patients with moderate and severe hepatic impairment, no specific dose recommendation can be given. **Paediatric population:** There is no relevant use of enfortumab vedotin in the paediatric population for the indication of locally advanced or metastatic urothelial cancer.

### Method of administration

Padcev is for intravenous use. The recommended dose must be administered by intravenous infusion over 30 minutes. Enfortumab vedotin must not be administered as an intravenous push or bolus injection. For instructions on reconstitution and dilution of the medicinal product before administration, see section 6.6 of the SPC. **Contraindications:** Hypersensitivity to the active substance or to any of the excipients listed in section 6.1 of the SPC. **Special warnings and precautions for use:**

**Traceability:** In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded. **Skin reactions:** Skin reactions are associated with enfortumab vedotin as a result of enfortumab vedotin binding to Nectin-4 expressed in the skin. Fever or flu-like symptoms may be the first sign of a severe skin reaction, and patients should be observed, if this occurs. Mild to moderate skin reactions, predominantly rash maculo-papular, have been reported with enfortumab vedotin. The incidence of skin reactions occurred at a higher rate when enfortumab vedotin was given in combination with pembrolizumab compared to enfortumab vedotin as monotherapy (see section 4.8 of the SPC). Severe cutaneous adverse reactions, including SJS and TEN, with fatal outcome have also occurred in patients treated with enfortumab vedotin, predominantly during the first cycle of treatment. Patients should be monitored starting with the first cycle and throughout treatment for skin reactions. Appropriate treatment such as topical corticosteroids and antihistamines can be considered for mild to moderate skin reactions. For suspected SJS or TEN, or in case of bullous lesions onset, withhold treatment immediately and refer to specialised care; histologic confirmation, including consideration of multiple biopsies, is critical to early recognition, as diagnosis and intervention can improve prognosis. Permanently discontinue Padcev for confirmed SJS or TEN, Grade 4 or recurrent Grade 3 skin reactions. For Grade 2 worsening, Grade 2 with fever or Grade 3 skin reactions, treatment should be withheld until Grade ≤1 and referral for specialised care should be considered. Treatment should be resumed at the same dose level or consider dose reduction by one dose level (see section 4.2 of the SPC). **Pneumonitis/ILD:** Severe, life-threatening or fatal pneumonitis/ILD have occurred in patients treated with enfortumab vedotin. The incidence of pneumonitis/ILD, including severe events occurred at a higher rate when enfortumab vedotin was given in combination with pembrolizumab compared to enfortumab vedotin as monotherapy (see section 4.8 of the SPC). Monitor patients for signs and symptoms indicative of pneumonitis/ILD such as hypoxia, cough, dyspnoea or interstitial infiltrates on radiologic exams. Corticosteroids should be administered for Grade ≥ 2 events (e.g., initial dose of 1–2 mg/kg/day prednisone or equivalent followed by a taper). Withhold Padcev for Grade 2 pneumonitis/ILD and consider dose reduction. Permanently discontinue Padcev for Grade ≥3 pneumonitis/ILD (see section 4.2 of the SPC). **Hyperglycaemia:** Hyperglycaemia and diabetic ketoacidosis (DKA), including fatal events, occurred in patients with and without pre-existing diabetes mellitus, treated with enfortumab vedotin (see section 4.8 of the SPC). Hyperglycaemia occurred more frequently in patients with pre-existing hyperglycaemia or a high body mass index (≥30 kg/m<sup>2</sup>). Patients treated with baseline HbA1c ≥8% were excluded from clinical studies. Blood glucose levels should be monitored prior to dosing and periodically throughout the course of treatment as clinically indicated in patients with or at risk for diabetes mellitus or hyperglycaemia. If blood glucose is elevated >13.9 mmol/L (>250 mg/dL), Padcev should be withheld until blood glucose is ≤13.9 mmol/L (<250 mg/dL) and treat as appropriate (see section 4.2 of the SPC). **Serious infections:** Serious infections such as sepsis (including fatal outcomes) have been reported in patients treated with Padcev. Patients should be carefully monitored during treatment for the emergence of possible serious infections. **Peripheral neuropathy:** Peripheral neuropathy, predominantly peripheral sensory neuropathy, has occurred with enfortumab vedotin, including Grade ≥3 reactions (see section 4.8 of the SPC). Patients with preexisting peripheral neuropathy Grade ≥2 were excluded from clinical studies. Patients should be monitored for symptoms of new or worsening peripheral neuropathy as these patients may require a delay, dose reduction or discontinuation of enfortumab vedotin (see Table 1). Padcev should be permanently discontinued for Grade ≥3 peripheral neuropathy (see section 4.2 of the SPC). **Ocular disorders:** Ocular disorders, predominantly dry eye, have occurred in patients treated with enfortumab vedotin (see section 4.8 of the SPC). Patients should be monitored for ocular disorders. Consider artificial tears for prophylaxis of dry eye and referral for ophthalmologic evaluation if ocular symptoms do not resolve or worsen. Infusion site extravasation: Skin and soft tissue injury following enfortumab vedotin administration has been observed when extravasation occurred (see section 4.8 of the SPC). Ensure good venous access prior to starting Padcev and monitor for possible infusion site extravasation during administration. If extravasation occurs, stop the infusion and monitor for adverse reactions. Embryo-fetal toxicity and contraception: Pregnant women should be informed of the potential risk to a fetus (see sections 4.6 and 5.3 of the SPC). Females of reproductive potential should be advised to have a pregnancy test within 7 days prior to starting treatment with enfortumab vedotin, to use effective contraception during treatment and for at least 6 months after stopping treatment. Men being treated with enfortumab vedotin are advised not to father a child during treatment and for at least 4 months following the last dose of Padcev. Patient information pack: The prescriber must discuss the risks of Padcev therapy, including combination therapy with pembrolizumab, with the patient. The patient should be provided with the patient information leaflet and patient card with each prescription. **Interactions:** Formal drug-drug interaction studies with enfortumab vedotin have not been conducted. Caution is advised in case of concomitant treatment with CYP3A4 inhibitors. Patients receiving concomitant strong CYP3A4 inhibitors should be monitored more closely for signs of toxicities. Strong CYP3A4 inducers may decrease the exposure of unconjugated MMAE with moderate effect (see section 5.2 of the SPC). **Undesirable effects:** Summary of the safety profile. **Enfortumab vedotin as monotherapy:** The safety of enfortumab vedotin was evaluated as monotherapy in 793 patients who received at least one dose of enfortumab vedotin 1.25 mg/kg in two phase 1 studies (EV-101 and EV-102), three phase 2 studies (EV-103, EV-201 and EV-203) and one phase 3 study (EV-301) (see Table 3). Patients were exposed to enfortumab vedotin for a median duration of 4.7 months (range: 0.3 to 55.7 months). The most common adverse reactions with enfortumab vedotin were alopecia (47.4%), decreased appetite (47.2%), fatigue (46.8%), diarrhoea (39.1%), peripheral sensory neuropathy (38.5%), nausea (37.8%), pruritus (33.4%), dysgeusia (30.4%), anaemia (29.1%), weight decreased (25.2%), rash maculo-papular (23.6%), dry skin (21.8%), vomiting (18.7%), aspartate aminotransferase increased (17%), hyperglycaemia (14.9%), dry eye (12.7%), alanine aminotransferase increased (12.7%) and rash (11.6%). The most common serious adverse reactions (≥2%) were diarrhoea (2.1%) and hyperglycaemia (2.1%). Twenty-one percent of patients permanently discontinued enfortumab vedotin for adverse reactions; the most common adverse reaction (≥2%) leading to dose discontinuation was peripheral sensory neuropathy (4.8%). Adverse reactions leading to dose interruption occurred in 62% of patients; the most common adverse reactions (≥2%) leading to dose interruption were peripheral sensory neuropathy (14.8%), fatigue (7.4%), rash maculo-papular (4%), aspartate aminotransferase increased (3.4%), alanine aminotransferase increased (3.2%), anaemia (3.2%), hyperglycaemia (3.2%), neutrophil count decreased (3%), diarrhoea (2.8%), rash (2.4%) and peripheral motor neuropathy (2.1%). Thirty-eight percent of patients required a dose reduction due to an adverse reaction; the most common adverse reactions (≥2%) leading to a dose reduction were peripheral sensory neuropathy (10.3%), fatigue (5.3%), rash maculo-papular (4.2%) and decreased appetite (2.1%). **Enfortumab vedotin in combination with pembrolizumab:** When enfortumab vedotin is administered in combination with pembrolizumab, refer to the SmPC for pembrolizumab prior to initiation of treatment. The safety of enfortumab vedotin was evaluated in combination with pembrolizumab in 564 patients who received at least one dose of enfortumab vedotin 1.25 mg/kg in combination with pembrolizumab in one phase 2 study (EV-103) and one phase 3 study (EV-302) (see Table 3). Patients were exposed to enfortumab vedotin in combination with pembrolizumab for a median duration of 9.4 months (range: 0.3 to 34.4 months). The most common adverse reactions with enfortumab vedotin in combination with pembrolizumab were peripheral sensory neuropathy (53.4%), pruritus (41.1%), fatigue (40.4%), diarrhoea (39.2%), alopecia (38.5%), rash maculo-papular (36%), weight decreased (36%), decreased appetite (33.9%), nausea (28.4%), anaemia (25.7%), dysgeusia (24.3%), dry skin (18.1%), alanine aminotransferase increased (16.8%), hyperglycaemia (16.7%), aspartate aminotransferase increased (15.4%), dry eye (14.4%), vomiting (13.3%), rash macular (11.3%), hypothyroidism (10.5%) and neutropenia (10.1%). The most common serious adverse reactions (≥2%) were diarrhoea (3%) and pneumonitis (2.3%). Thirty-six percent of patients permanently discontinued enfortumab vedotin for adverse reactions; the most common adverse reactions (≥2%) leading to discontinuation were peripheral sensory neuropathy (12.2%) and rash maculo-papular (2%). Adverse reactions leading to dose interruption of enfortumab vedotin occurred in 72% of patients. The most common adverse reactions (≥2%) leading to dose interruption were peripheral sensory neuropathy (17%), rash maculo-papular (6.9%), diarrhoea (4.8%), fatigue (3.7%), pneumonitis (3.7%), hyperglycaemia (3.4%), neutropenia (3.2%), alanine aminotransferase increased (3%), pruritus (2.3%) and anaemia (2%). Adverse reactions leading to dose reduction of enfortumab vedotin occurred in 42.4% of patients. The most

common adverse reactions (≥2%) leading to dose reduction were peripheral sensory neuropathy (9.9%), rash maculo-papular (6.4%), fatigue (3.2%), diarrhoea (2.3%) and neutropenia (2.1%). **Tabulated summary of adverse reactions:** Adverse reactions observed during clinical studies of enfortumab vedotin as monotherapy or in combination with pembrolizumab, or reported from post-marketing use of enfortumab vedotin are listed in this section by frequency category. Frequency categories are defined as follows: very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥1/10,000 to <1/1,000); very rare (<1/10,000); not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

**Table 3. Adverse reactions in patients treated with enfortumab vedotin**

	Monotherapy	In combination with pembrolizumab
<b>Infections and infestations</b>		
Common	Sepsis	Sepsis
<b>Blood and lymphatic system disorders</b>		
Very common	Anaemia	Anaemia
Not known <sup>1</sup>	Neutropenia, febrile neutropenia, neutrophil count decreased	Neutropenia, febrile neutropenia, neutrophil count decreased
<b>Endocrine disorders</b>		
Very common		Hypothyroidism
<b>Metabolism and nutrition disorders</b>		
Very common	Hyperglycaemia, decreased appetite	Hyperglycaemia, decreased appetite
Not known <sup>1</sup>	Diabetic ketoacidosis	Diabetic ketoacidosis
<b>Nervous system disorders</b>		
Very common	Peripheral sensory neuropathy, dysgeusia	Peripheral sensory neuropathy, dysgeusia
Common	Neuropathy peripheral, peripheral motor neuropathy, peripheral sensorimotor neuropathy, paraesthesia, hypoaesthesia, gait disturbance, muscular weakness	Peripheral motor neuropathy, peripheral sensorimotor neuropathy, paraesthesia, hypoaesthesia, gait disturbance, muscular weakness
Uncommon	Demyelinating polyneuropathy, polyneuropathy, neurotoxicity, motor dysfunction, dysaesthesia, muscle atrophy, neuralgia, peroneal nerve palsy, sensory loss, skin burning sensation, burning sensation	Neurotoxicity, dysaesthesia, myasthenia gravis, neuralgia, peroneal nerve palsy, skin burning sensation
<b>Eye disorders</b>		
Very common	Dry eye	Dry eye
<b>Respiratory, thoracic, and mediastinal disorders</b>		
Very common		Pneumonitis/ILD <sup>2</sup>
Common	Pneumonitis/ILD <sup>2</sup>	
<b>Gastrointestinal disorders</b>		
Very common	Diarrhoea, vomiting, nausea	Diarrhoea, vomiting, nausea
<b>Skin and subcutaneous tissue disorders</b>		
Very common	Alopecia, pruritus, rash, rash maculo-papular, dry skin	Alopecia, pruritus, rash, rash maculo-papular, dry skin, rash macular
Common	Drug eruption, skin exfoliation, conjunctivitis, dermatitis bullous, blister, stomatitis, palmar-plantar erythrodysesthesia syndrome, eczema, erythema, rash erythematous, rash macular, rash papular, rash pruritic, rash vesicular	Rash, skin exfoliation, conjunctivitis, dermatitis bullous, blister, stomatitis, palmar-plantar erythrodysesthesia syndrome, eczema, erythema, rash erythematous, rash papular, rash pruritic, rash vesicular, erythema multiforme, dermatitis
Uncommon	Dermatitis exfoliative generalised, erythema multiforme, exfoliative rash, pemphigoid, rash maculovesicular, dermatitis, dermatitis allergic, dermatitis contact, intertrigo, skin irritation, stasis dermatitis, bull blister	Drug eruption, dermatitis exfoliative generalised, exfoliative rash, pemphigoid, dermatitis contact, intertrigo, skin irritation, stasis dermatitis
Not known <sup>1</sup>	Toxic epidermal necrolysis, skin hyperpigmentation, skin discoloration, pigmentation disorder, Stevens Johnson syndrome, epidermal necrosis, symmetrical drug-related intertriginous and flexural exanthema	Toxic epidermal necrolysis, skin hyperpigmentation, skin discoloration, pigmentation disorder, Stevens Johnson syndrome, epidermal necrosis, symmetrical drug-related intertriginous and flexural exanthema
<b>Musculoskeletal and connective tissue disorders</b>		
Common		Myositis
<b>General disorders and administration site conditions</b>		
Very common	Fatigue	Fatigue
Common	Infusion site extravasation	Infusion site extravasation
<b>Investigations</b>		
Very common	Alanine aminotransferase increased, aspartate aminotransferase increased, weight decreased	Alanine aminotransferase increased, aspartate aminotransferase increased, weight decreased
Common		Lipase increased
<b>Injury, poisoning and procedural complications</b>		
Common	Infusion related reaction	Infusion related reaction

<sup>1</sup>Based on global post-marketing experience.

<sup>2</sup>Includes: acute respiratory distress syndrome, autoimmune lung disease, immune-mediated lung disease, interstitial lung disease, lung opacity, organising pneumonia, pneumonitis, pulmonary fibrosis, pulmonary toxicity and sarcoidosis. **Description of selected adverse reactions: Immunogenicity:** A total of 697 patients were tested for immunogenicity to enfortumab vedotin 1.25 mg/kg as monotherapy; 16 patients were confirmed to be positive at baseline for anti-drug antibody (ADA), and in patients that were negative at baseline (N=681), a total of 24 (3.5%) were positive post baseline. A total of 490 patients were tested for immunogenicity against enfortumab vedotin following enfortumab vedotin in combination with pembrolizumab; 24 patients were confirmed to be positive at baseline for ADA, and in patients that were negative at baseline (N=466), a total of

14 (3%) were positive post baseline. The incidence of treatment-emergent anti-enfortumab vedotin antibody formation was consistent when assessed following enfortumab vedotin administration as monotherapy and in combination with pembrolizumab. Due to the limited number of patients with antibodies against Padcev, no conclusions can be drawn concerning a potential effect of immunogenicity on efficacy, safety or pharmacokinetics. **Skin reactions:** In clinical studies of enfortumab vedotin as monotherapy, skin reactions occurred in 57% (452) of the 793 patients treated with enfortumab vedotin 1.25 mg/kg. Severe (Grade 3 or 4) skin reactions occurred in 14% (108) of patients and a majority of these reactions included rash maculo-papular, stomatitis, rash erythematous, rash or drug eruption. The median time to onset of severe skin reactions was 0.7 months (range: 0.1 to 8.2 months). Serious skin reactions occurred in 4.3% (34) of patients. Of the patients who experienced skin reactions and had data regarding resolution (N=366), 61% had complete resolution, 24% had partial improvement, and 15% had no improvement at the time of their last evaluation. Of the 39% of patients with residual skin reactions at last evaluation, 38% had Grade ≥2 events. In clinical studies of enfortumab vedotin in combination with pembrolizumab, skin reactions occurred in 70% (392) of the 564 patients and a majority of these skin reactions included rash maculo-papular, rash macular and rash papular. Severe (Grade 3 or 4) skin reactions occurred in 17% (97) of patients (Grade 3: 16%, Grade 4: 1%). The median time to onset of severe skin reactions was 1.7 months (range: 0.1 to 17.2 months). Of the patients who experienced skin reactions and had data regarding resolution (N=391), 59% had complete resolution, 30% had partial improvement, and 10% had no improvement at the time of their last evaluation. Of the 41% of patients with residual skin reactions at last evaluation, 27% had Grade ≥2 events. **Pneumonitis/ILD:** In clinical studies of enfortumab vedotin as monotherapy, pneumonitis/ILD occurred in 26 (3.3%) of the 793 patients treated with enfortumab vedotin 1.25 mg/kg. Less than 1% of patients experienced severe (Grade 3 or 4) pneumonitis/ILD (Grade 3: 0.5%, Grade 4: 0.3%). Pneumonitis/ILD led to discontinuation of enfortumab vedotin in 0.5% of patients. There were no deaths from pneumonitis/ILD. The median time to onset of any grade pneumonitis/ILD was 2.7 months (range: 0.6 to 6.0 months) and the median duration for pneumonitis/ILD was 1.6 months (range: 0.1 to 43.0 months). Of the 26 patients who experienced pneumonitis/ILD, 8 (30.8%) had resolution of symptoms. In clinical studies of enfortumab vedotin in combination with pembrolizumab, pneumonitis/ILD occurred in 58 (10.3%) of the 564 patients. Severe (Grade 3 or 4) pneumonitis/ILD occurred in 20 patients (Grade 3: 3.0%, Grade 4: 0.5%). Pneumonitis/ILD led to discontinuation of enfortumab vedotin in 2.1% of patients. Two patients experienced a fatal event of pneumonitis/ILD. The median time to onset of any grade pneumonitis/ILD was 4 months (range: 0.3 to 26.2 months). **Hyperglycaemia:** In clinical studies of enfortumab vedotin as monotherapy, hyperglycaemia (blood glucose >13.9 mmol/L) occurred in 17% (133) of the 793 patients treated with enfortumab vedotin 1.25 mg/kg. Serious events of hyperglycaemia occurred in 2.5% of patients, 7% of patients developed severe (Grade 3 or 4) hyperglycaemia and 0.3% of patients experienced fatal events, one event each of hyperglycaemia and diabetic ketoacidosis. The incidence of Grade 3-4 hyperglycaemia increased consistently in patients with higher body mass index and in patients with higher baseline haemoglobin A1C (HbA1c). The median time to onset of hyperglycaemia was 0.5 months (range: 0 to 20.3). Of the patients who experienced hyperglycaemia and had data regarding resolution (N=106), 66% had complete resolution, 19% had partial improvement, and 15% had no improvement at the time of their last evaluation. Of the 34% of patients with residual hyperglycaemia at last evaluation, 64% had Grade ≥2 events. **Peripheral neuropathy:** In clinical studies of enfortumab vedotin as monotherapy, peripheral neuropathy occurred in 53% (422) of the 793 patients treated with enfortumab vedotin 1.25 mg/kg. Five percent of patients experienced severe (Grade 3 or 4) peripheral neuropathy including sensory and motor events. The median time to onset of Grade ≥2 peripheral neuropathy was 5 months (range: 0.1 to 20.2). Of the patients who experienced neuropathy and had data regarding resolution (N=340), 14% had complete resolution, 46% had partial improvement, and 41% had no improvement at the time of their last evaluation. Of the 86% of patients with residual neuropathy at last evaluation, 51% had Grade ≥2 events. **Ocular disorders:** In clinical studies of enfortumab vedotin as monotherapy, 30% of patients experienced dry eye during treatment with enfortumab vedotin 1.25 mg/kg. Treatment was interrupted in 1.5% of patients and 0.1% of patients permanently discontinued treatment due to dry eye. Severe (Grade 3) dry eye only occurred in 3 patients (0.4%). The median time to onset of dry eye was 1.7 months (range: 0 to 30.6 months). **Special populations: Elderly:** Enfortumab vedotin in combination with pembrolizumab has been studied in 173 patients <65 years and 391 patients ≥65 years. Generally, adverse event frequencies were higher in patients ≥65 years of age compared to <65 years of age, particularly for serious adverse events (56.3%, and 35.3%, respectively) and Grade ≥3 events (80.3% and 64.2%, respectively), similar to observations with the chemotherapy comparator. **Overdose:** There is no known antidote for overdose with enfortumab vedotin. In case of overdose, the patient should be closely monitored for adverse reactions, and supportive treatment should be administered as appropriate taking into consideration the half-life of 3.6 days (ADC) and 2.6 days (MMAE).

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

**België/Belgique:** Federal Agentschap voor Geneesmiddelen en Gezondheidsproducten / Agence fédérale des médicaments et des produits de santé; [www.fagg.be](http://www.fagg.be) / [www.afmps.be](http://www.afmps.be); Afdeling Vigilantie / Division Vigilance; Website/Site internet: [www.eunijwerkingmelden.be](http://www.eunijwerkingmelden.be) / [www.notifierunneffettidesirable.be](http://www.notifierunneffettidesirable.be); e-mail: [adr@fagg-afmps.be](mailto:adr@fagg-afmps.be)

**Ireland:** HPRA Pharmacovigilance, Website: [www.hpra.ie](http://www.hpra.ie) or Astellas Pharma Co. Ltd. Tel.: +353 1 467 1555, E-mail: [irishdrugsafety@astellas.com](mailto:irishdrugsafety@astellas.com).

**Nederland:** Nederlands Bijwerkingen Centrum Lareb; Website: [www.lareb.nl](http://www.lareb.nl)

**Luxembourg/Luxemburg :** Centre Régional de Pharmacovigilance de Nancy ou Division de la pharmacie et des médicaments de la Direction de la santé / Site internet : [www.guichet.lu/pharmacovigilance](http://www.guichet.lu/pharmacovigilance)

### MARKETING AUTHORISATION HOLDER:

Astellas Pharma Europe B.V. Sylviusweg 62, 2333 BE Leiden, The Netherlands

**MARKETING AUTHORISATION NUMBERS:** EU/1/21/1615/001 & EU/1/21/1615/002

**DATE OF REVISION OF THE TEXT:** December 2024

**Job Bag Number:** MAT-BX-PAD-2025-00004

Detailed information on this medicinal product is available on the website of the European Medicines Agency <http://www.ema.europa.eu>.

**Ireland:** Astellas Pharma Co., Ltd., Tel.: +353 1 467 1555. SPC may be found at [www.medicines.ie](http://www.medicines.ie).

Delivery Status: subject to medical prescription.

Astellas Pharma B.V.

NL: Sylviusweg 62, 2333BE Leiden, Netherlands

BE/LU: Mediaalane 50, 1800 Vilvoorde, Belgium

IE: Legal classification: S1A.



**Prescribing Information: PADCEV™ (enfortumab vedotin) 20 mg and 30 mg powder for concentrate for solution for infusion**

For full prescribing information refer to the Summary of Product Characteristics (SPC).

**Presentation:** One vial of PADCEV powder for concentrate for solution for infusion contains either 20 mg or 30 mg enfortumab vedotin. After reconstitution, each mL of solution contains 10 mg of enfortumab vedotin. Enfortumab vedotin is comprised of a fully human IgG1 kappa antibody, conjugated to the microtubule-disrupting agent monomethyl auristatin E (MMAE) via a protease-cleavable maleimidocaproyl valine-citrulline linker.

**Indications:** PADCEV, in combination with pembrolizumab, is indicated for the first-line treatment of adult patients with unresectable or metastatic urothelial cancer who are eligible for platinum-containing chemotherapy. PADCEV as monotherapy is indicated for the treatment of adult patients with locally advanced or metastatic urothelial cancer who have previously received a platinum-containing chemotherapy and a programmed death receptor-1 or programmed death-ligand 1 inhibitor (see section 5.1 of the SPC).

**Posology and method of administration:** Treatment with PADCEV should be initiated and supervised by a physician experienced in the use of anti-cancer therapies. PADCEV is for intravenous use. It must not be administered as an intravenous push or bolus injection. Good venous access prior to starting treatment should be ensured (see section 4.4 of the SPC). As monotherapy, the recommended dose of enfortumab vedotin is 1.25 mg/kg (up to a maximum of 125 mg for patients >100 kg). It must be administered as an intravenous infusion over 30 minutes on Days 1, 8 and 15 of a 28-day cycle until disease progression or unacceptable toxicity. When given in combination with pembrolizumab, the recommended dose of enfortumab vedotin is 1.25 mg/kg (up to a maximum of 125 mg for patients >100 kg) administered as an intravenous infusion over 30 minutes on Days 1 and 8 of every 3-week (21-day) cycle until disease progression or unacceptable toxicity. The recommended dose of pembrolizumab is either 200 mg every 3 weeks or 400 mg every 6 weeks administered as an intravenous infusion over 30 minutes. Patients should be administered pembrolizumab after enfortumab vedotin when given on the same day. Refer to the pembrolizumab SmPC for additional dosing information of pembrolizumab. For information on recommended dose reductions of enfortumab vedotin for adverse reactions as well as instructions on dose modifications (interruption, reduction and discontinuation) in patients experiencing adverse reactions refer to section 4.2 of the SPC. **Special Populations: Elderly:** No dose adjustment is necessary in patients >65 years of age (see section 5.2 of the SPC). **Renal impairment:** No dose adjustment is necessary in patients with mild [creatinine clearance (CrCL) >60-90 mL/min], moderate (CrCL 30–60 mL/min) or severe (CrCL 15–<30 mL/min) renal impairment. Enfortumab vedotin has not been evaluated in patients with end stage renal disease (CrCL <15 mL/min) (see section 5.2 of the SPC). **Hepatic impairment:** No dose adjustment is necessary in patients with mild hepatic impairment [total bilirubin of 1 to 1.5 x upper limit of normal (ULN) and aspartate transaminase (AST) any, or total bilirubin ≤ ULN and AST > ULN]. Enfortumab vedotin has only been evaluated in a limited number of patients with moderate and severe hepatic impairment. Hepatic impairment is expected to increase the systemic exposure to MMAE (the cytotoxic drug); therefore, patients should be closely monitored for potential adverse events. Due to the sparsity of the data in patients with moderate and severe hepatic impairment, no specific dose recommendation can be given (see section 5.2 of the SPC). **Paediatric population:** There is no relevant use of enfortumab vedotin in the paediatric population for the indication of locally advanced or metastatic urothelial cancer.

**Contraindications:** Hypersensitivity to the active substance or to any of the excipients listed in section 6.1 of the SPC.

**Special warnings and precautions for use: Traceability:** In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded. **Skin reactions:** Skin reactions are associated with enfortumab vedotin as a result of enfortumab vedotin binding to Nectin-4 expressed in the skin. Fever or flu-like symptoms may be the first sign of a severe skin reaction, and patients should be observed, if this occurs. Mild to moderate skin reactions, predominantly rash maculo-papular, have been reported with enfortumab vedotin. The incidence of skin reactions occurred at a higher rate when enfortumab vedotin was given in combination with pembrolizumab compared to enfortumab vedotin as monotherapy (see section 4.8 of the SPC). Severe cutaneous adverse reactions, including Stevens-Johnson syndrome (SJS) and Toxic Epidermal Necrolysis (TEN), with fatal outcome have also occurred in patients treated with enfortumab vedotin, predominantly during the first cycle of treatment. Patients should be monitored starting with the first cycle and throughout treatment for skin reactions. Appropriate treatment such as topical corticosteroids and antihistamines can be considered for mild to moderate skin reactions. For suspected SJS or TEN, or in case of bullous lesions onset, withhold treatment immediately and refer to specialised care; histologic confirmation, including consideration of multiple biopsies, is critical to early recognition, as diagnosis and intervention can improve prognosis. Permanently discontinue PADCEV for confirmed SJS or TEN, Grade 4 or recurrent Grade 3 skin reactions. For Grade 2 worsening, Grade 2 with fever or Grade 3 skin reactions, treatment should be withheld until Grade ≤1 and referral for specialised care should be considered. Treatment should be resumed at the same dose level or consider dose reduction by one dose level (see section 4.2 of the SPC). **Pneumonitis/Interstitial Lung Disease (ILD):** Severe, life-threatening or fatal pneumonitis/ILD have occurred in patients treated with enfortumab vedotin. The incidence of pneumonitis/ILD, including severe events occurred at a higher rate when enfortumab vedotin was given in combination with pembrolizumab compared to enfortumab vedotin as monotherapy (see section 4.8 of the SPC). Monitor patients for signs and symptoms indicative of pneumonitis/ILD such as hypoxia, cough, dyspnoea or interstitial infiltrates on radiologic exams. Corticosteroids should be administered for Grade ≥2 events (e.g., initial dose of 1–2 mg/kg/day prednisone or equivalent followed by a taper). Withhold PADCEV for Grade 2 pneumonitis/ILD and consider dose reduction. Permanently discontinue PADCEV for Grade ≥3 pneumonitis/ILD (see section 4.2 of the SPC). **Hyperglycaemia:** Hyperglycaemia and diabetic ketoacidosis (DKA), including fatal events, occurred in patients with and without pre-existing diabetes mellitus, treated with enfortumab vedotin (see section 4.8 of the SPC). Hyperglycaemia occurred more frequently in patients with pre-existing hyperglycaemia or a high body mass index (≥30 kg/m<sup>2</sup>). Patients with baseline HbA1c ≥8% were excluded from clinical studies. Blood glucose levels should be monitored prior to dosing and periodically throughout the course of treatment as clinically indicated in patients with or at risk for diabetes mellitus or hyperglycaemia. If blood glucose is elevated >13.9 mmol/L

(>250 mg/dL), PADCEV should be withheld until blood glucose is ≤13.9 mmol/L (<250 mg/dL) and treat as appropriate (see section 4.2 of the SPC). **Serious infections:** Serious infections such as sepsis (including fatal outcomes) have been reported in patients treated with PADCEV. Patients should be carefully monitored during treatment for the emergence of possible serious infections. **Peripheral neuropathy:** Peripheral neuropathy, predominantly peripheral sensory neuropathy, has occurred with enfortumab vedotin, including Grade ≥3 reactions (see section 4.8 of the SPC). Patients with pre-existing peripheral neuropathy Grade ≥2 were excluded from clinical studies. Patients should be monitored for symptoms of new or worsening peripheral neuropathy as these patients may require a delay, dose reduction or discontinuation of enfortumab vedotin. PADCEV should be permanently discontinued for Grade ≥3 peripheral neuropathy (see section 4.2 of the SPC). **Ocular disorders:** Ocular disorders, predominantly dry eye, have occurred in patients treated with enfortumab vedotin (see section 4.8 of the SPC). Patients should be monitored for ocular disorders. Consider artificial tears for prophylaxis of dry eye and referral for ophthalmologic evaluation if ocular symptoms do not resolve or worsen. **Infusion site extravasation:** Skin and soft tissue injury following enfortumab vedotin administration has been observed when extravasation occurred (see section 4.8 of the SPC). Ensure good venous access prior to starting PADCEV and monitor for possible infusion site extravasation during administration. If extravasation occurs, stop the infusion and monitor for adverse reactions. **Embryo-fœtal toxicity and contraception:** Pregnant women should be informed of the potential risk to a foetus (see sections 4.6 and 5.3 of the SPC). Females of reproductive potential should be advised to have a pregnancy test within 7 days prior to starting treatment with enfortumab vedotin, to use effective contraception during treatment and for at least 6 months after stopping treatment. Men being treated with enfortumab vedotin are advised not to father a child during treatment and for at least 4 months following the last dose of PADCEV. **Patient information pack:** The prescriber must discuss the risks of PADCEV therapy, including combination therapy with pembrolizumab, with the patient. The patient should be provided with the patient information leaflet and patient card with each prescription.

**Effects on ability to drive and use machines:** PADCEV has no or negligible influence on the ability to drive and use machines.

**Interactions:** Formal drug-drug interaction studies with enfortumab vedotin have not been conducted. Caution is advised in case of concomitant treatment with CYP3A4 inhibitors. Patients receiving concomitant strong CYP3A4 inhibitors (e.g. bupropion, clarithromycin, cobicistat, indinavir, itraconazole, nefazodone, nelfinavir, posaconazole, ritonavir, saquinavir, telaprevir, telithromycin, voriconazole) should be monitored more closely for signs of toxicity. Strong CYP3A4 inducers (e.g. rifampicin, carbamazepine, phenobarbital, phenytoin, St John's wort [*Hypericum perforatum*]) may decrease the exposure of unconjugated MMAE with moderate effect (see section 5.2 of the SPC).

**Fertility, pregnancy and lactation: Women of childbearing potential/ Contraception in males and females:** Refer to 'Special warnings and precautions for use' section above. **Pregnancy:** PADCEV can cause foetal harm when administered to pregnant women based upon findings from animal studies. PADCEV is not recommended during pregnancy and in women of childbearing potential not using effective contraception. **Breast-feeding:** Breast-feeding should be discontinued during PADCEV treatment and for at least 6 months after the last dose. **Fertility:** Men being treated with this medicinal product are advised to have sperm samples frozen and stored before treatment. There are no data on the effect of PADCEV on human fertility.

**Undesirable effects: Summary of the safety profile: Enfortumab vedotin as monotherapy:** The safety of enfortumab vedotin was evaluated as monotherapy in 793 patients who received at least one dose of enfortumab vedotin 1.25 mg/kg in two phase 1 studies (EV-101 and EV-102), three phase 2 studies (EV-103, EV-201 and EV-203) and one phase 3 study (EV-301) (see Table 3 in section 4.8 of the SPC). Patients were exposed to enfortumab vedotin for a median duration of 4.7 months (range: 0.3 to 55.7 months). The most common adverse reactions with enfortumab vedotin were alopecia (47.7%), decreased appetite (47.2%), fatigue (46.8%), diarrhoea (39.1%), peripheral sensory neuropathy (38.5%), nausea (37.8%), pruritus (33.4%), dysgeusia (30.4%), anaemia (29.1%), weight decreased (25.2%), rash maculo-papular (23.6%), dry skin (21.8%), vomiting (18.7%), aspartate aminotransferase increased (17%), hyperglycaemia (14.9%), dry eye (12.7%), alanine aminotransferase increased (12.7%) and rash (11.6%). The most common serious adverse reactions (≥2%) were diarrhoea (2.1%) and hyperglycaemia (2.1%). Twenty-one percent of patients permanently discontinued enfortumab vedotin for adverse reactions; the most common adverse reaction (≥2%) leading to dose discontinuation was peripheral sensory neuropathy (4.8%). Adverse reactions leading to dose interruption occurred in 62% of patients; the most common adverse reactions (≥2%) leading to dose interruption were peripheral sensory neuropathy (14.8%), fatigue (7.4%), rash maculo-papular (4%), aspartate aminotransferase increased (3.4%), alanine aminotransferase increased (3.2%), anaemia (3.2%), hyperglycaemia (3.2%), neutrophil count decreased (3%), diarrhoea (2.8%), rash (2.4%) and peripheral motor neuropathy (2.1%). Thirty-eight percent of patients required a dose reduction due to an adverse reaction; the most common adverse reactions (≥2%) leading to a dose reduction were peripheral sensory neuropathy (10.3%), fatigue (5.3%), rash maculo-papular (4.2%) and decreased appetite (2.1%). *Enfortumab vedotin in combination with pembrolizumab:* When enfortumab vedotin is administered in combination with pembrolizumab, refer to the SPC for pembrolizumab prior to initiation of treatment. The safety of enfortumab vedotin was evaluated in combination with pembrolizumab in 564 patients who received at least one dose of enfortumab vedotin 1.25 mg/kg in combination with pembrolizumab in one phase 2 study (EV-103) and one phase 3 study (EV-302) (see Table 3). Patients were exposed to enfortumab vedotin in combination with pembrolizumab for a median duration of 9.4 months (range: 0.3 to 34.4 months). The most common adverse reactions with enfortumab vedotin in combination with pembrolizumab were peripheral sensory neuropathy (53.4%), pruritus (41.1%), fatigue (40.4%), diarrhoea (39.2%), alopecia (38.5%), rash maculo-papular (36%), weight decreased (36%), decreased appetite (33.9%), nausea (28.4%), anaemia (25.7%), dysgeusia (24.3%), dry skin (18.1%), alanine aminotransferase increased (16.8%), hyperglycaemia (16.7%), aspartate aminotransferase increased (15.4%), dry eye (14.4%), vomiting (13.3%), rash macular (11.3%), hypothyroidism (10.5%) and neutropenia (10.1%). The most common serious adverse reactions (≥2%) were diarrhoea (3%) and pneumonitis (2.3%). Thirty-six percent of patients permanently discontinued enfortumab vedotin for adverse reactions; the most common adverse reactions (≥2%) leading to discontinuation were peripheral sensory neuropathy (12.2%) and rash

maculo-papular (2%). Adverse reactions leading to dose interruption of enfortumab vedotin occurred in 72% of patients. The most common adverse reactions (≥2%) leading to dose interruption were peripheral sensory neuropathy (17%), rash maculo-papular (6.9%), diarrhoea (4.8%), fatigue (3.7%), pneumonitis (3.7%), hyperglycaemia (3.4%), neutropenia (3.2%), alanine aminotransferase increased (3%), pruritus (2.3%) and anaemia (2%). Adverse reactions leading to dose reduction of enfortumab vedotin occurred in 42.4% of patients. The most common adverse reactions (≥2%) leading to dose reduction were peripheral sensory neuropathy (9.9%), rash maculo-papular (6.4%), fatigue (3.2%), diarrhoea (2.3%) and neutropenia (2.1%). **Summary of adverse reactions:** Adverse reactions observed during clinical studies of enfortumab vedotin as monotherapy or in combination with pembrolizumab, or reported from post-marketing use of enfortumab vedotin are listed in this section according to Medical Dictionary for Regulatory Activities (MedDRA) system organ classification by frequency category. Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness. Frequency categories are defined as follows: very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥1/10,000 to <1/1,000); very rare (<1/10,000); not known (cannot be estimated from the available data). **Infections and infestations:** (monotherapy and in combination with pembrolizumab) Common: Sepsis. **Blood and lymphatic system disorders:** (monotherapy and in combination with pembrolizumab) Very common: Anaemia. Not known<sup>1</sup>: Neutropenia, febrile neutropenia, neutrophil count decreased. **Endocrine disorders:** (in combination with pembrolizumab) Very common: Hypothyroidism. **Metabolism and nutrition disorders:** (monotherapy and in combination with pembrolizumab) Very common: Hyperglycaemia, decreased appetite. Not known<sup>1</sup>: Diabetic ketoacidosis. **Nervous system disorders:** (monotherapy and in combination with pembrolizumab) Very common: Peripheral sensory neuropathy, dysgeusia. (monotherapy) Common: Neuropathy peripheral, peripheral motor neuropathy, peripheral sensorimotor neuropathy, paraesthesia, hypoaesthesia, gait disturbance, muscular weakness. (in combination with pembrolizumab) Common: Peripheral motor neuropathy, peripheral sensorimotor neuropathy, paraesthesia, hypoaesthesia, gait disturbance, muscular weakness. (monotherapy) Uncommon: Demyelinating polyneuropathy, polyneuropathy, neurotoxicity, motor dysfunction, dysaesthesia, muscle atrophy, neuralgia, peroneal nerve palsy, sensory loss, skin burning sensation, burning sensation. (in combination with pembrolizumab) Uncommon: Neurotoxicity, dysaesthesia, myasthenia gravis, neuralgia, peroneal nerve palsy, skin burning sensation. **Eye disorders:** (monotherapy and in combination with pembrolizumab) Very common: Dry eye. **Respiratory, thoracic, and mediastinal disorders:** (in combination with pembrolizumab) Very common: Pneumonitis/ILD<sup>2</sup>. (monotherapy) Common: Pneumonitis/ILD<sup>2</sup>. **Gastrointestinal disorders:** (monotherapy and in combination with pembrolizumab) Very common: Diarrhoea, vomiting, nausea. **Skin and subcutaneous tissue disorders:** (monotherapy) Very common: Alopecia, pruritus, rash, rash maculo-papular, dry skin. (in combination with pembrolizumab) Very common: Alopecia, pruritus, rash maculo-papular, dry skin, rash macular. (monotherapy) Common: Drug eruption, skin exfoliation, conjunctivitis, dermatitis bullous, blister, stomatitis, palmar-plantar erythrodysesthesia syndrome, eczema, erythema, rash erythematous, rash macular, rash papular, rash pruritic, rash vesicular. (in combination with pembrolizumab) Common: Rash, skin exfoliation, conjunctivitis, dermatitis bullous, blister, stomatitis, palmar-plantar erythrodysesthesia syndrome, eczema, erythema, rash erythematous, rash papular, rash pruritic, rash vesicular, erythema multiforme, dermatitis. (monotherapy) Uncommon: Dermatitis exfoliative generalised, erythema multiforme, exfoliative rash, pemphigoid, rash maculovesicular, dermatitis, dermatitis allergic, dermatitis contact, intertrigo, skin irritation, stasis dermatitis, blood blister. (in combination with pembrolizumab) Uncommon: Drug eruption, dermatitis exfoliative generalised, exfoliative rash, pemphigoid, dermatitis contact, intertrigo, skin irritation, stasis dermatitis. (monotherapy and in combination with pembrolizumab) Not known<sup>1</sup>: TEN, SJS, epidermal necrosis, skin hyperpigmentation, skin discoloration, pigmentation disorder, symmetrical drug-related intertriginous and flexural exanthema. **Musculoskeletal and connective tissue disorders:** (in combination with pembrolizumab) Common: Myositis. **General disorders and administration site conditions:** (monotherapy and in combination with pembrolizumab) Very common: Fatigue. (monotherapy and in combination with pembrolizumab) Common: Infusion site extravasation. **Investigations:** (monotherapy and in combination with pembrolizumab) Very common: Alanine aminotransferase increased, aspartate aminotransferase increased, weight decreased. (in combination with pembrolizumab) Common: Lipase increased. **Injury, poisoning and procedural complications:** (monotherapy and in combination with pembrolizumab) Common: Infusion related reaction.

<sup>1</sup>Based on global post-marketing experience.

<sup>2</sup>Includes: acute respiratory distress syndrome, autoimmune lung disease, immune-mediated lung disease, interstitial lung disease, lung opacity, organizing pneumonia, pneumonitis, pulmonary fibrosis, pulmonary toxicity and sarcoidosis.

**Description of selected adverse reactions, Immunogenicity:** A total of 697 patients were tested for immunogenicity to enfortumab vedotin 1.25 mg/kg as monotherapy; 16 patients were confirmed to be positive at baseline for anti-drug antibody (ADA), and in patients that were negative at baseline (N=681), a total of 24 (3.5%) were positive post baseline. A total of 490 patients were tested for immunogenicity against enfortumab vedotin following enfortumab vedotin in combination with pembrolizumab; 24 patients were confirmed to be positive at baseline for ADA, and in patients that were negative at baseline (N=466), a total of 14 (3%) were positive post baseline. The incidence of treatment-emergent anti-enfortumab vedotin antibody formation was consistent when assessed following enfortumab vedotin administration as monotherapy and in combination with pembrolizumab. Due to the limited number of patients with antibodies against PADCEV, no conclusions can be drawn concerning a potential effect of immunogenicity on efficacy, safety or pharmacokinetics. **Skin reactions:** In clinical studies of enfortumab vedotin as monotherapy, skin reactions occurred in 57% (452) of the 793 patients treated with enfortumab vedotin 1.25 mg/kg. Severe (Grade 3 or 4) skin reactions occurred in 14% (108) of patients and a majority of these reactions included rash maculo-papular, stomatitis, rash erythematous, rash or drug eruption. The median time to onset of severe skin reactions was 0.7 months (range: 0.1 to 8.2 months). Serious skin reactions occurred in 4.3% (34) of patients. Of the patients who experienced skin reactions and had data regarding resolution (N=366), 61% had complete resolution, 24% had

partial improvement, and 15% had no improvement at the time of their last evaluation. Of the 39% of patients with residual skin reactions at last evaluation, 38% had Grade ≥2 events. In clinical studies of enfortumab vedotin in combination with pembrolizumab, skin reactions occurred in 70% (392) of the 564 patients and a majority of these skin reactions included rash maculo-papular, rash macular and rash papular. Severe (Grade 3 or 4) skin reactions occurred in 17% (97) of patients (Grade 3: 16%, Grade 4: 1%). The median time to onset of severe skin reactions was 1.7 months (range: 0.1 to 17.2 months). Of the patients who experienced skin reactions and had data regarding resolution (N=391), 59% had complete resolution, 30% had partial improvement, and 10% had no improvement at the time of their last evaluation. Of the 41% of patients with residual skin reactions at last evaluation, 27% had Grade ≥2 events. **Pneumonitis/ILD:** In clinical studies of enfortumab vedotin as monotherapy, pneumonitis/ILD occurred in 26 (3.3%) of the 793 patients treated with enfortumab vedotin 1.25 mg/kg. Less than 1% of patients experienced severe (Grade 3 or 4) pneumonitis/ILD (Grade 3: 0.5%, Grade 4: 0.3%). Pneumonitis/ILD led to discontinuation of enfortumab vedotin in 0.5% of patients. There were no deaths from pneumonitis/ILD. The median time to onset of any grade pneumonitis/ILD was 2.7 months (range: 0.6 to 6.0 months) and the median duration for pneumonitis/ILD was 1.6 months (range: 0.1 to 43.0 months). Of the 26 patients who experienced pneumonitis/ILD, 8 (30.8%) had resolution of symptoms. In clinical studies of enfortumab vedotin in combination with pembrolizumab, pneumonitis/ILD occurred in 58 (10.3%) of the 564 patients. Severe (Grade 3 or 4) pneumonitis/ILD occurred in 20 patients (Grade 3: 3.0%, Grade 4: 0.5%). Pneumonitis/ILD led to discontinuation of enfortumab vedotin in 2.1% of patients. Two patients experienced a fatal event of pneumonitis/ILD. The median time to onset of any grade pneumonitis/ILD was 4 months (range: 0.3 to 26.2 months). **Hyperglycaemia:** In clinical studies of enfortumab vedotin as monotherapy, hyperglycaemia (blood glucose >13.9 mmol/L) occurred in 17% (133) of the 793 patients treated with enfortumab vedotin 1.25 mg/kg. Serious events of hyperglycaemia occurred in 2.5% of patients, 7% of patients developed severe (Grade 3 or 4) hyperglycaemia and 0.3% of patients experienced fatal events, one event each of hyperglycaemia and diabetic ketoacidosis. The incidence of Grade 3–4 hyperglycaemia increased consistently in patients with higher body mass index and in patients with higher baseline haemoglobin A1C (HbA1c). The median time to onset of hyperglycaemia was 0.5 months (range: 0 to 20.3). Of the patients who experienced hyperglycaemia and had data regarding resolution (N=106), 66% had complete resolution, 19% had partial improvement, and 15% had no improvement at the time of their last evaluation. Of the 34% of patients with residual hyperglycaemia at last evaluation, 64% had Grade ≥2 events. **Peripheral neuropathy:** In clinical studies of enfortumab vedotin as monotherapy, peripheral neuropathy occurred in 53% (422) of the 793 patients treated with enfortumab vedotin 1.25 mg/kg. Five percent of patients experienced severe (Grade 3 or 4) peripheral neuropathy including sensory and motor events. The median time to onset of Grade ≥2 peripheral neuropathy was 5 months (range: 0.1 to 20.2). Of the patients who experienced neuropathy and had data regarding resolution (N=340), 14% had complete resolution, 46% had partial improvement, and 41% had no improvement at the time of their last evaluation. Of the 86% of patients with residual neuropathy at last evaluation, 51% had Grade ≥2 events. **Ocular disorders:** In clinical studies of enfortumab vedotin as monotherapy, 30% of patients experienced dry eye during treatment with enfortumab vedotin 1.25 mg/kg. Treatment was interrupted in 1.5% of patients and 0.1% of patients permanently discontinued treatment due to dry eye. Severe (Grade 3) dry eye only occurred in 3 patients (0.4%). The median time to onset of dry eye was 1.7 months (range: 0 to 30.6 months). **Special populations:** Elderly: Enfortumab vedotin in combination with pembrolizumab has been studied in 173 patients <65 years and 391 patients ≥65 years. Generally, adverse event frequencies were higher in patients ≥65 years of age compared to <65 years of age, particularly for serious adverse events (56.3%, and 35.3%, respectively) and Grade ≥3 events (80.3% and 64.2%, respectively), similar to observations with the chemotherapy comparator. Prescribers should consult the full SPC in relation to other adverse reactions.

**Overdose:** There is no known antidote for overdose with enfortumab vedotin. In case of overdose, the patient should be closely monitored for adverse reactions, and supportive treatment should be administered as appropriate taking into consideration the half-life of 3.6 days (ADC and 2.6 days (MMAE)).

**Cost (excluding VAT):** PADCEV 20 mg powder for concentrate for solution for infusion x 1 vial: £578  
PADCEV 30 mg powder for concentrate for solution for infusion x 1 vial: £587

**Legal classification:** POM

**Marketing Authorisation numbers:**

PADCEV 20 mg powder for concentrate for solution for infusion PLGB 00166/0432.

PADCEV 30 mg powder for concentrate for solution for infusion PLGB 00166/0433.

**Marketing Authorisation Holder:**

Astellas Pharma Ltd, 300 Dashwood Lang Road, Bourne Business Park, Addlestone, United Kingdom, KT15 2NX.

**Date of Preparation of Prescribing Information:** February 2025

**Job Bag Number:** MAT-GB-PAD-2025-00017

**Further information available from:** Astellas Pharma Ltd, Medical Information 0800 783 5018.

For full prescribing information, refer to the SPC, which may be found at: <https://www.medicines.org.uk/emc>.

**Adverse events should be reported. Reporting forms and information can be found at [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. Adverse events should also be reported to Astellas Pharma Ltd. on 0800 783 5018.**

*The hyperlinks on this page will take you to non-Astellas websites. Astellas does not endorse or accept liability for sites controlled by third-parties.*